



09/889687  
PCT/AU00/00025

REC'D 29 FEB 2000	
1/250	PCT

Patent Office  
Canberra

4

I, LEANNE MYNOTT, TEAM LEADER EXAMINATION SUPPORT AND SALES hereby certify that annexed is a true copy of the Provisional specification in connection with Application No. PP 8230 for a patent by ALCHEMIA PTY LTD filed on 18 January 1999.



WITNESS my hand this  
Twenty-second day of February 2000

A handwritten signature in dark ink, appearing to read 'L. Mynott'.

LEANNE MYNOTT  
TEAM LEADER EXAMINATION  
SUPPORT AND SALES

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

AUSTRALIA  
Patents Act 1990

PROVISIONAL SPECIFICATION

**Applicant(s):** ALCHEMIA PTY LTD  
A.C.N. 071 666 334

**Invention Title:** PROTECTING GROUPS FOR CARBOHYDRATE SYNTHESIS

The invention is described in the following statement:

## PROTECTING GROUPS FOR CARBOHYDRATE SYNTHESIS

This invention relates to methods of synthesis of glycoconjugates, and in particular to orthogonally protected carbohydrate building blocks. The invention provides collections of orthogonally protected monosaccharides as universal building blocks for the synthesis of; glycoconjugates of non-carbohydrate molecules, neo-glycoconjugates and oligosaccharides. This orthogonal protection strategy allows for the specific deprotection of any substituent on the saccharide ring, and greatly facilitates targeted or library focused carbohydrate related syntheses .

### 15 BACKGROUND OF THE INVENTION

Oligosaccharides are important components of a variety of different types of biological molecules, and are involved in antigenic recognition and cell-cell interactions. In many cases, bio-molecules require conjugation with a carbohydrate component in order to be fully functional. In order to enable investigation of the biological function, and to exploit the exquisite biochemical and antigenic specificity of oligosaccharides, it is essential to have access to highly defined, specific synthetic oligosaccharides. Therefore achieving efficient, cost-effective synthesis of oligosaccharides and glycoconjugates by either solution or solid phase methods is of the utmost importance.

This task is enormously complicated by the complexity of oligosaccharides. Because of the number of sites which can carry substituents, and the number of possible ways in which two saccharide molecules can be linked, the number of permutations is enormously high.

In naturally-occurring oligosaccharides D-glucose, D-galactose L-fucose, D-mannose, D-glucosamine and D-galactosamine are among the most common sugar residues. To construct oligosaccharides and carbohydrate conjugates

using these sugars, current methodologies require long, protracted syntheses, involving synthesis of as many as one hundred different specially-protected sugar donors in order to cover adequately all the possible permutations of glycosidic link formation (eg. 1-3, 1-4), link type (eg.  $\alpha$  or  $\beta$ ) and to include all possible branching points in the oligosaccharide.

Previously orthogonal protection of bi-functional molecules has been a widely used technique in organic chemistry which provided general building blocks for selected syntheses. But orthogonal protection in the case of molecules with a greater degree of functionalisation is quite rare. Our technology involves penta-functional monosaccharide building blocks, which require a much higher level of chemical specificity to attain the appropriate orthogonality.

Orthogonal protection has been defined by Merrifield as follows:

"The principle of orthogonal stability requires that only those protecting functions should be used that can be cleaved under different reaction conditions without affecting the other functions present"  
(Merrifield, 1977)

Although the use of orthogonal protection would greatly facilitate carbohydrate related synthesis, there has been limited success in devising suitable protecting groups and methods.

Wong synthesised a universal building block with chloroacetyl, *p*-methoxybenzyl, levulinyl and *tert*-butyldiphenylsilyl protecting groups selectively removable with sodium bicarbonate, trifluoroacetic acid, hydrazine and hydrogen fluoride-pyridine respectively, on a galactopyranose ring with an aryl-thio leaving group at the glycosidic position. This building block was used solely to synthesise a 6-hexanate glycoside. The subsequent recombinant oligosaccharide library formation focused on using the 6-hexanate derivatised building block which

exhibits only four degrees of orthogonality (Wong *et al*, 1998).

Similarly Kunz and coworkers synthesised an orthogonally protected D-glucopyranose derivative, but  
5 synthetic manipulations were only performed on the aglycon. These authors describe orthogonal protection of hydroxyl groups on a monosaccharide linked at C2 via a thioglycoside group to a solid support or to a succinimide moiety. In  
10 this case the protecting groups are acetyl or methyl at C1, allyl at C4, ethoxyethyl at C5, and tert-butyldiphenylsilyl at C6. The thioglycoside anchor functionalized in the side-chain is stated to be crucial. Again there is no suggestion that this protection system can be used for substituted sugars. The Kunz's orthogonally protected  
15 building block was not used for glycosylation or construction of glycoconjugates or neo-glycoconjugates, by directly attaching functionalities to the pyranose ring (Wunberg *et al*. 1998).


In our earlier International Patent Applications  
20 No. PCT/AU97/00544, No. PCT/AU98/00131 and No. PCT/AU98/00808, we described protecting and linking groups which enabled oligosaccharides and aminooligosaccharides to be synthesised using solid phase methods of the type which for many years have applied to  
25 peptide synthesis. In addition the protecting groups, described therein were useful for solution-phase synthesis. The entire disclosures of these specifications are incorporated herein by this reference.

We have now devised new types of building blocks  
30 which greatly facilitate the synthesis of oligosaccharides and glycoconjugates, using orthogonally-protected saccharide building blocks with five degrees of orthogonality. These building blocks contain a leaving group or latent leaving group at the glycosidic position,  
35 and another four orthogonally protected functional groups around the carbohydrate ring.


Using our approach with six universal building blocks based on six of the most common naturally occurring sugars, any one of the one hundred sugars referred to above may be quickly synthesised in a facile manner, using  
5 simple, well-known protecting group chemistry. The years of work and complex protection strategies required to produce these one hundred building blocks by previously-available methods can be avoided by use of our six universal building blocks, which do not require a high level of skill to use,  
10 and enable one to achieve the synthesis of a specific desired oligosaccharide or glycoconjugate much faster and more efficiently than previously possible.

#### SUMMARY OF THE INVENTION

15 In its most general aspect the invention provides a universal monosaccharide building block of General Formula I or General Formula II



I



II

20

in which

A is a leaving group, including but not limited to groups such as -SR, where R is alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl,  
25 cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, halogen, trichloroacetimidoyl-, sulphoxide, acyl, azido, thiocyanate or -O-alkenyl;

X is hydrogen, O, N or N<sub>3</sub>;

X<sub>1</sub> is hydrogen, -CH<sub>2</sub>O-, -CH<sub>2</sub>NH-, -CH<sub>3</sub>, -CH<sub>2</sub>N<sub>3</sub> or  
30 -COO-; and

B, C, D and E are any protecting groups that can be cleaved orthogonally.

It will be appreciated that as a consequence of stoichiometry and valence bond theory B, C, D and E are absent when X is hydrogen or N<sub>3</sub> and E is absent when X<sub>1</sub> is hydrogen, CH<sub>3</sub> or N<sub>3</sub>.

5           The following non-limiting sets have been designated as orthogonal to each other on the basis of their cleavage conditions. A protecting group is classified in a particular set according to its lability to the cleavage conditions for a particular set and its  
10 stability to the cleavage conditions required for the removal of those groups in the remaining sets. Each set is to be taken to include, but is not be limited, by the members thereof.

          Of the sets defined, set 1, the 'Base Solvolysis' set, is of particular importance, because in addition to  
15 the fact that the members of this set are considered to be orthogonal to the members of the remaining sets, some members of this set are also considered to be orthogonal to each other. Where this is the case the alternative  
20 condition of cleavage that provides orthogonality is specified in brackets following the listing of the protecting group.

1. Base Solvolysis  
25       a) for hydroxy protection:

          acyl-type protecting groups, eg. chloroacetate  
          (also thiourea-sensitive)  
          bromoacetate (also pyridine-sensitive)  
30       carbonates, eg. Alloc (Pd<sup>0</sup>)  
          Fmoc (β-elimination)  
          Troc  
          p-nitrophenylsulphonylethyloxy carbonyl)  
          levanoyl (also hydrazine sensitive)  
35

b) for amino protection:

Dde (primary amine-sensitive)  
tetraphthaloyl  
5 dichlorophthaloyl  
2,5-dimethyl-pyrroyl (primary amine-sensitive)  
benzyloxycarbonyl  
pentenyl

10 2. Fluoride Ion Sensitive  
for hydroxy protection:

t-butyldiphenylsilyl  
triisopropylsilyl  
15 trimethylsilylethyl  
triphenylsilylethyl  
(all cleavable with HF/Pyridine)

20 3. Reduction-Sensitive

trifluoromethyl  
trichloromethyloxymethyl  
trichloromethyloxycarbonate  
(all cleavable with zinc/acetic acid)

25 4.  $\beta$ -Elimination Sensitive Base Labile Protecting Groups

ethoxyethyl  
cyanoethyl  
30 NSC (p-nitrobenzyl-sulphonylethyloxycarbonyl)  
p-nitrobenzyl-sulphonylethyl

5. Hydrogenolysis-Sensitive Protecting Groups

35 naphthylmethyl  
substituted naphthylmethyl

6. Oxidation-Sensitive Protecting Groups:

- 5           p-methoxybenzyl  
          3,4-dimethoxybenzyl  
          2,4,6-trimethoxybenzyl  
          piperinoyl  
          acylamidobenzyl  
          azidobenzyl  
10          p-azido-m-chlorobenzyl

7. Allylic Protecting Groups

Cleavable with Pd<sup>0</sup> complexes


15 8. Photolabile Protecting Groups:

- o-nitrobenzyloxycarbonate  
          o-nitrobenzyl  
          dinitrobenzyl  
20          2-oxo-1,2-diphenylethyl

\*9. Protecting Groups Removable by Relay Deprotection

- 25          methylthioethyl  
          acyloxybenzyl  
          benzylthioethyl.

30           In one preferred embodiment, the invention  
          provides a compound of General Formula III



III


in which

A, X and X<sub>1</sub> are as defined for General Formulae I and II, and

5 B<sub>1</sub>, C<sub>1</sub>, D<sub>1</sub> and E<sub>1</sub> are orthogonal carbohydrate protecting groups (ie. an orthogonal set) selected from protecting group sets 1, 2, 6 and 8.

Another preferred embodiment provides a compound of General Formula IV

10




in which

15 A, X and X<sub>1</sub> are as defined for General Formulae I and II, and

B<sub>2</sub>, C<sub>2</sub>, D<sub>2</sub> and E<sub>2</sub> are selected from the members of protecting group set 1, and in themselves constitute an orthogonal set, for example the carbohydrate-protecting groups levanoyl (ammonia-labile), chloroacetate (thiourea-labile), *p*-methoxybenzyloxycarbonyl (oxidation-labile) and 20 2-trimethylsilylethylcarbonate (fluoride ion-labile).

This embodiment provides universal building blocks with protecting groups selected from the protecting groups of set 1.

25 In a third preferred embodiment the invention provides a compound of General Formula V



in which

A, X and  $X_1$  are as defined for General Formula I and II, and

5         $B_3$ ,  $C_3$ ,  $D_3$  and  $E_3$  are an orthogonal set of protecting groups selected from amongst the members of set 1 and from the remaining orthogonal sets.

      This embodiment provides orthogonally protected building blocks, the protecting group constituents of which  
10       may be selected from within set 1 and from the remaining sets.

      It will be clearly understood that the invention is not limited to use with monosaccharides, but is also applicable to any compound in which substituents are linked  
15       to a pyranose or furanose ring, such as sugar analogues.

      For the purposes of this specification it will be clearly understood that the word "comprising" means "including but not limited to", and that the word "comprises" has a corresponding meaning.

20       For the purposes of this specification "orthogonal cleavage" is defined as the regioselective cleavage of a hydroxy or amino protecting group from a carbohydrate, in which the cleavage conditions do not compromise the stability of the other protecting or  
25       functional groups on the molecule. Such cleavages can be effected in any order of priority. "Cleaved orthogonally" and "orthogonal cleavage" are taken to be synonymous.

#### DETAILED DESCRIPTION OF THE INVENTION

30       Abbreviations used herein are as follows:

Alloc	Allyloxycarbonyl
Bn	Benzyl
Bu	Butyl
35 DCM	Dichloromethane
Dde	N-1-(4,4-Dimethyl-2,6-dioxocyclohexylidene)ethyl


	Dde-OH	6-Hydroxy-6-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl
	DMAP	<i>N,N'</i> -Dimethylaminopyridine
	DMF	<i>N,N'</i> -Dimethylformamide
5	DMTST	Dimethyl(methylthio)sulphoniumtrifluoromethanesulphonate
	EEDQ	1-isobutyloxycarbonyl-2-isobutyloxy-1,2-dihydroquinoline
	EtOAc	Ethyl acetate
10	EtOH	Ethanol
	FAB-MS	Fast atom bombardment mass spectrometry
	HRMS	High resolution mass spectrometry
	Fmoc	Fluoromethoxycarbonyl
	MBHA	Methyl benzyhydramine resin
15	Me	Methyl
	MeOH	Methanol
	NCS	<i>p</i> -Nitrobenzyl-sulphonylethyloxycarbonyl
	NMR	Nuclear magnetic resonance
	ODmab	4-{ <i>N</i> -[1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-3-methylbutyl]-amino}benzyl alcohol
20		
	PEG	Polyethylene glycol
	tBu	Tertiary-butyl
	TFA	Trifluoroacetic acid
	THF	Tetrahydrofuran
25	Troc	2,2,2-Trichloroethoxycarbonyl

The invention provides universal building blocks which are useful in the solution and solid phase synthesis of oligosaccharides. The reaction scheme for synthesis of each target molecule is designed so as to specify the orthogonally-protected functional groups which must be freed for glycosylation, and those which need to be capped with a protecting group such as benzyl, benzoyl or another such group which remains uncleaved until the end of the synthesis, to avoid competition during glycosylations later in the synthesis.

When participation during the glycosylation reaction is required, the 2-hydroxyl is selectively deprotected and reprotected with a benzoyl group which, again, remains until the completion of the synthesis. In  
5 the case of 2-deoxy 2-aminosugars, if participation or stereoselectivity is required the non-participating Dde group is removed and replaced with a tetrachlorophthaloyl or 2,5-dimethylpyrrole group.

10 Example 1      Synthesis of an Exemplary Tetrasaccharide

A strategy for synthesis of the tetrasaccharide of formula VI is set out in Scheme 1.




VI



In solution phase, protecting groups A and C from the first sugar residue of the target molecule (residue [4]) are selectively removed, and the sites capped by a permanent protecting group, eg. benzoyl group. The residue  
5 is then coupled to the resin, followed by selective removal of protecting group B. In solution phase, protecting group A from sugar residue [3] is selectively removed, and the site is capped by a permanent protecting group. Residue [3] is then linked to the resin-bound sugar residue via a  
10 glycosylation reaction. Protecting group C from the new disaccharide is removed, and residue [2] is linked via a glycosylation. Protecting group A is finally selectively removed to regenerate the 6-hydroxyl group which is linked with residue 1.

15  
It will be apparent to the person skilled in the art that while the invention has been described in some detail for the purposes of clarity and understanding, various modifications and alterations to the embodiments  
20 and methods described herein may be made without departing from the scope of the inventive concept disclosed in this specification.

Example 2      Synthesis of a Fully Orthogonally Protected  
Thioglycoside Building Block, methyl 6-O-(t-  
butyldiphenylsilyl)-3-O-(p-chlorobenzoyl)-2-  
deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-  
ylidene)ethylamino]-4-O-tetrahydropyranyl-1-  
thio-β-D glucopyranoside



10      **1**      **Methyl 4,6-O-benzylidene-2-deoxy-2-[1-(4,4-dimethyl-**  
**2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio-β-D**  
**glucopyranoside**

A mixture of methyl 2-deoxy-2-[1-(4,4-dimethyl-  
2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio-β-D  
15 glucopyranoside (20g, 54mmol), α,α-dimethoxytoluene  
(9.78 g, 64 mmol) and p-toluenesulphonic acid (50 mg) in  
dry acetonitrile (100 ml), was stirred at 60°C for 2 hours.  
The reaction mixture was cooled to room temperature and  
adjusted to pH 7 with the addition of triethylamine. The  
20 solvent was removed *in vacuo* and the residue taken up in  
CH<sub>2</sub>Cl<sub>2</sub> (200 ml), washed with brine (50 ml), with water

(50 ml) and dried over  $\text{MgSO}_4$ . The organic phase was concentrated to give a yellow solid, methyl 4,6-O-benzylidene-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside (24.5 g, 98%).

**2 Methyl 4,6-O-benzylidene-3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside**

A mixture of methyl 4,6-O-Benzylidene-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside (6.3 g, 13.5 mmol), p-chlorobenzoylchloride (2.6 ml, 20 mmol) and 4-dimethylaminopyridine (2.44 g, 40 mmol) in dry 1,2-dichloroethane (100 ml), was stirred at room temperature overnight. The resultant suspension was filtered, the filtrate diluted with chloroform (100 ml) and washed with diluted brine (3 x 50 ml,  $\text{H}_2\text{O}$ /Brine, 2/1). The organic phase was dried over  $\text{MgSO}_4$  and the solvent removed in vacuo to give yellow solid. The residue chromatographed EtOAc/Hexane 1:1 as the mobile phase to give methyl 4,6-O-benzylidene-3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside (6.4 g, 80%).

**3 Methyl 3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside**

A mixture of methyl 4,6-O-benzylidene-3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside (2.51 g, 4.20 mmol) and 50% aqueous solution of tetrafluoroboric acid (1 ml) in acetonitrile

(25 ml), was stirred at room temperature for 2 hours. The pH was adjusted to 7 with the addition of triethylamine and the resultant suspension concentrated. The residue was crystallised from diisopropyl ether-ethyl acetate to give  
5 methyl 3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside (1.7 g, 79%).

4     **Methyl 6-O-(t-butyldiphenylsilyl)-3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside**  
10

A mixture of methyl 3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)-ethylamino]-1-thio- $\beta$ -D glucopyranoside (1.00 g, 1.95 mmol),  
15 t-butyldiphenylsilylchloride (536 mg, 1.95) and 4-dimethylaminopyridine (238 mg, 1.95 mmol), in 1,2-dichloroethane (30 ml), was stirred under reflux for 6 hour. The reaction mixture was cooled to room  
20 temperature, diluted with chloroform (60 ml) and washed with diluted brine (3 x 50 ml, brine/water, 1:2), dried over magnesiumsulphate. The solvent was removed *in vacuo* to and the chromatographed using hexane/ethylacetate 1:1 as the mobile phase to give a white solid, methyl 6-O-(t-  
25 butyldiphenylsilyl)-3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside (1.1 g, 75%)

5    **Methyl 6-O-(t-butyldiphenylsilyl)-3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-4-O-tetrahydropyranyl-1-thio- $\beta$ -D glucopyranoside**

5            A mixture of methyl 6-O-(t-butyldiphenylsilyl)-3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-1-thio- $\beta$ -D glucopyranoside (500 mg, 0.6 mmol), 3,4-dihydro-2H-pyran (5 ml) and p-toluenesulphonic acid (5 mg) in dry  
10    acetonitrile (10 ml) was stirred at room temperature for 1 hour. The reaction mixture was adjusted to pH 7 with the addition of triethylamine and then evaporated to dryness. The residue was taken up in dichloromethane (30 ml), washed with water (2 x 10 ml) and the organic phase dried over  
15     $\text{MgSO}_4$ . The solvent was removed in vacuo and the residue chromatographed using hexane/EtOAc 2:1 as the mobile phase to give methyl 6-O-(t-butyldiphenylsilyl)-3-O-(p-chlorobenzoyl)-2-deoxy-2-[1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethylamino]-4-O-tetrahydropyranyl-  
20    1-thio- $\beta$ -D glucopyranoside (420 mg, 85%).

References cited herein are listed on the following pages, and are incorporated herein by this reference.

REFERENCES

Henke, S.

Angew. Chem. Int. Ed., 1998 37 2503-2505

5

Merrifield, R. B.

Pept., Proc. Am. Pept. Symp., 5<sup>th</sup>, 1977 488.

Wong, C-H, Ye, X-S and Zhang, Z.

10 J. Am. Chem. Soc., 1998 120 7137-7138.

ALCHEMIA PTY LTD

18 January 1999